

1

2

6

8

2

2

2

2

3

2

2

2.

2.

1 10. The method of claim 9, wherein the functional effect is measured
2 by determining changes in the electrical activity of cells expressing the G-protein alpha
3 subunit polypeptide.

1 11. The method of claim 10, wherein the changes in electrical activity
2 are measured by an assay selected from the group consisting of a voltage clamp assay, a
3 patch clamp assay, a radiolabeled ion flux assay, or a fluorescence assay using voltage
4 sensitive dyes.

1 12. The method of claim 9, wherein the functional effect is determined
2 by measuring changes in the level of phosphorylation of sensory cell specific proteins.

1 13. The method of claim 9, wherein the functional effect is determined
2 by measuring changes in transcription levels of sensory cell specific genes.

1 14. The method of claim 9, wherein the functional effect is determined
2 by measuring changes in intracellular cAMP, cGMP, IP₃, DAG, or Ca²⁺.

1 15. The method of claim 14, wherein the changes in intracellular
2 cAMP or cGMP are measured using immunoassays.

1 16. The method of claim 9, wherein the cell or cell membrane is
2 attached to a solid substrate.

1 17. The method of claim 9, wherein the cell is a eukaryotic cell.

1 18. The method of claim 17, wherein the cell is a human cell.

1 19. The method of claim 18, wherein the cell is an HEK 293 cell.

1 20. The method of claim 9, wherein the G-protein alpha subunit
2 polypeptide is co-expressed with GPCR-B3 or GPCR-B4.

1 21. The method of claim 1, wherein the G-protein alpha subunit
2 polypeptide is linked to a solid phase.

1 22. The method of claim 21, wherein the G-protein alpha subunit
2 polypeptide is covalently linked to the solid phase.

009210-820460

(i) expressing a sensory cell specific G-protein alpha subunit polypeptide in an HEK 293 host cell, wherein the G-protein alpha subunit polypeptide comprises greater than 70% amino acid sequence identity to a polypeptide having a sequence of SEQ ID NO:2;

9 (iii) contacting the host cell with the compound that modulates
10 sensory signaling in sensory cells; and

1 24. The method of claim 23, wherein the sensory cell specific G-
2 protein coupled receptor is GPCR-B3 or GPCR-B4.